

## MUTUAL INJECTIVE HULLS

S. K. JAIN, S. R. LÓPEZ-PERMOUTH AND S. RAZA SYED

**ABSTRACT.** The mutual injective hull of an arbitrary family of modules is constructed. Applications to the calculations of quasi-injective and  $\pi$ -injective hulls of direct sums are given.

**1. Introduction.** Consider the direct sum  $M = \bigoplus_{i \in I} M_i$  of a family of right modules  $\{M_i \mid i \in I\}$ . Assume, for simplicity, that  $I$  is finite or  $R$  is right noetherian. If each  $M_i$  is injective, then  $M$  is injective. However, if each  $M_i$  is merely quasi-injective (or  $\pi$ -injective) it does not necessarily follow that  $M$  is quasi-injective (resp.  $\pi$ -injective). In either case,  $M$  is quasi-injective ( $\pi$ -injective) if and only if the  $M_i$ 's are "mutually injective" in the sense that whenever  $j \neq k$ ,  $M_j$  is  $M_k$ -injective. If the  $M_i$ 's are arbitrary, the question then arises to determine formulae for the quasi-injective hull  $q(M)$  and the  $\pi$ -injective hull  $\pi(M)$  of  $M$  as a direct sum of essential extensions of the  $M_i$ 's. These essential extensions should then be quasi-injective ( $\pi$ -injective) and mutually injective. We start by focusing on the problem of producing, for an arbitrary family of modules  $\{N_j \mid j \in J\}$ , over an arbitrary ring  $R$  a family of "smallest" essential extensions  $\{\tilde{N}_j \mid j \in J\}$  which are mutually injective, namely a *mutual injective hull* of  $\{N_j \mid j \in J\}$ .

Theorem 3.1 proves the existence and Theorem 3.2 provides formulae for the mutual injective hulls of arbitrary families of modules. The construction of mutual injective hulls allows us then to show that if  $I$  is finite or  $R$  is right noetherian

$$q(M) = \bigoplus_{i \in I} q(\tilde{M}_i) = \bigoplus_{i \in I} q(\tilde{M}_i)$$

and

$$\pi(M) = \bigoplus_{i \in I} \pi(\tilde{M}_i) = \bigoplus_{i \in I} \pi(\tilde{M}_i).$$

**2. Definitions and preliminaries.** A family of right modules  $\{N_j \mid j \in J\}$  over an arbitrary ring  $R$  is said to be *mutually injective* if, for all  $j \neq k$  in  $J$ ,  $N_j$  is  $N_k$ -injective. Given a family  $\{N_j \mid j \in J\}$  with corresponding injective hulls  $\{E(N_j) \mid j \in J\}$ , a family  $\{\tilde{N}_j \mid j \in J\}$  is said to be a *mutual injective hull* for  $\{N_j \mid j \in J\}$  (in  $\{E(N_j) \mid j \in J\}$ ) if

- (i) for all  $j \in J$ ,  $N_j \subseteq \tilde{N}_j \subseteq E(N_j)$ ,
- (ii)  $\{\tilde{N}_k \mid k \in J\}$  is mutually injective, and

---

Research supported by NSF Grant No. INT9210491 and Baker Fund Award, Ohio University, first author. Received by the editors June 12, 1995.

AMS subject classification: 16D50.

© Canadian Mathematical Society 1996.

- (iii) for any mutually injective family  $\{N'_k \mid k \in J\}$  if for all  $j \in J$ ,  $N_j \subseteq N'_j \subseteq E(N_j)$ , then for all  $j \in J$ ,  $\tilde{N}_j \subseteq N'_j$ .

Throughout this paper, all modules are right and unital unless otherwise stated. For an  $R$ -module  $M$ , the symbols  $E(M)$  (or  $\hat{M}$ ),  $q(M)$  and  $\pi(M)$  will denote, respectively, the injective hull, quasi-injective hull and  $\pi$ -injective hull of  $M$ . An  $R$ -module  $M$  is injective relative to an  $R$ -module  $N$  (or  $N$ -injective) if

$$\text{hom}_R(N, E(M))N \subseteq M.$$

$M$  is quasi-injective if  $M$  is injective relative to itself, i.e.,  $M$  is invariant under  $\text{End}(E(M))$ .  $M$  is  $\pi$ -injective if for any two submodules  $A, B \subset M$  with  $A \cap B = 0$ , each of the projection maps  $A \oplus B \rightarrow A$  and  $A \oplus B \rightarrow B$  can be extended to an endomorphism of  $M$ . Equivalently,  $M$  is  $\pi$ -injective iff for each idempotent  $e \in \text{End}(E(M))$ ,  $eM \subset M$ . It is well known that a module is  $\pi$ -injective if and only if it is quasi-continuous. For any two modules  $M_1$  and  $M_2$ ,  $M_1 \subset' M_2$  will denote that  $M_1$  is essential in  $M_2$ .

We record below some well-known lemmas which are used throughout the paper.

LEMMA 2.1. *The quasi-injective hull of  $M$  is given by  $q(M) = \text{End}(E(M))M$  and the  $\pi$ -injective hull of  $M$  is given by  $\pi(M) = SM$ , where  $S$  is the subring generated by the idempotent elements of  $\text{End}(E(M))$ .*

LEMMA 2.2.  *$M$  is  $\pi$ -injective if and only if  $E(M) = \bigoplus_{i \in I} M_i$  implies  $M = \bigoplus \Sigma(M_i \cap M)$ .*

PROOF. See ([3], Theorem 1.1).

LEMMA 2.3. (a) *If  $M$  is  $N$ -injective, then  $M$  is both  $K$ - and  $(N/K)$ -injective for each submodule  $K$  of  $N$ .*

(b) *If  $M$  is  $N_i$ -injective,  $i \in I$ , then  $M$  is  $\bigoplus_{i \in I} N_i$ -injective.*

(c) *If  $M$  is  $N$ -injective and  $M \subset' M'$  is  $N$ -injective, then  $M'$  is  $N$ -injective.*

(d) *If  $M$  is  $N$ -injective, then  $M$  is  $SN$ -injective for any subring  $S$  of  $\text{End}(E(N))$ . In particular,  $M$  is  $\pi(N)$ - (and  $q(N)$ -) injective.*

PROOF. It is straightforward. (See, for example, ([1] Proposition 16.13, p. 188).

LEMMA 2.4. *Let  $M$  and  $N$  be  $R$ -modules. Let  $M_1 = \text{hom}_R(E(N), E(M))N + M$ . Then:*

(i)  *$M_1$  is  $N$ -injective;*

(ii) *if  $M'$  is any other  $R$ -module such that  $M \subset M' \subset E(M)$  and  $M'$  is  $N$ -injective, then  $M_1 \subset M'$ .*

PROOF. Follows immediately from the definition of  $N$ -injectivity.

We call  $M_1$  (in Lemma 2.4) the  $N$ -injective hull of  $M$  (in  $E(M)$ ), and denote it by  $E_N(M)$ .

LEMMA 2.5. *If  $M = \bigoplus_{i \in I} M_i$  is  $\pi$ -injective, then  $M_i$  is  $\pi$ -injective and  $\{M_i \mid i \in J\}$  is mutually injective. In case  $I$  is finite or  $R$  is noetherian, the converse also holds.*

PROOF. See [4].

**3. Main results.**

**THEOREM 3.1.** (Existence). *Every family of modules  $\{N_j \mid j \in J\}$  over an arbitrary ring  $R$  with corresponding injective hulls  $\{E(N_j) \mid j \in J\}$  has a mutual injective hull.*

**PROOF.** Using the relative injective hulls introduced in Lemma 2.4, we define the sequence

$$N_i^{(0)} = N_i, \\ N_i^{(k)} = E_{\sum_{j \neq i} N_j^{(k-1)}}(N_i^{(k-1)}), \quad k = 1, 2, 3, \dots$$

Let  $\tilde{N}_i = \cup N_i^{(k)}$ . We show  $\tilde{N}_i$  is  $\tilde{N}_j$ -injective for  $i \neq j$ . So, let  $f: \tilde{N}_j \rightarrow E(\tilde{N}_i)$  be an  $R$ -homomorphism. By our construction,  $N_i^{(k)}$  is  $\sum_{j \neq i} N_j^{(k-1)}$ -injective hence  $N_j^{(k-1)}$ -injective for  $j \neq i$ . Also,  $E(\tilde{N}_i) = E(N_i)$ . So  $f(\tilde{N}_j) \subset \cup_k N_i^{(k)} = \tilde{N}_i, j \neq i$ . Thus  $\tilde{N}_i$  is  $\tilde{N}_j$ -injective.

Let  $\{N'_i\}$  be a mutually injective family such that  $N_i \subset N'_i \subset \hat{N}_i$ , for all  $i$ . We show that  $\tilde{N}_i \subset N'_i$  for all  $i$ . Now  $N_i^{(1)}$  is relative  $\sum_{j \neq i} N_j$ -injective hull of  $N_i$  in  $E(N_i)$ , and since  $N'_i$  is also  $\sum_{j \neq i} N_j$ -injective, it follows that

$$N_i^{(1)} \subset N'_i, \text{ for all } i.$$

So  $N'_i$  is  $\sum_{j \neq i} N_j^{(1)}$ -injective, yielding

$$N_i^{(2)} \subset N'_i.$$

Proceeding like this, we obtain  $N_i^{(k)} \subset N'_i \forall k$ ; hence  $\tilde{N}_i \subset N'_i$  as desired.

The family  $(\tilde{N}_i)_{i \in I}$  of  $R$ -modules obtained in the above theorem will be called the *mutual injective hull of the family  $(N_i)_{i \in I}$  in  $(E(N_i))_{i \in I}$ .*

The next theorem shows that the sequence of  $R$ -modules  $\{N_i^{(k)}\}, k = 1, 2, \dots$  constructed above to obtain  $\tilde{N}_i$  indeed becomes stationary after the second step, i.e.,  $N_i^{(2)} = N_i^{(k)}$  for all  $k \geq 2$ .

**THEOREM 3.2.** *With the notation as in the proof of Theorem 3.1,  $\tilde{N}_i = N_i^{(2)}$ . Indeed,*

$$\tilde{N}_i = N_i + \sum_{j \neq i} \text{hom}(\hat{N}_j, \hat{N}_i)N_j + \sum_{j \neq i} \text{hom}(\hat{N}_j, \hat{N}_i) \text{hom}(\hat{N}_i, \hat{N}_j)N_i,$$

where  $\hat{N} = E(N)$ .

**PROOF.** For simplicity we will consider a family with only two modules  $N_1$  and  $N_2$ . The general case works similarly.

$$N_1^{(1)} = \text{hom}(\hat{N}_2, \hat{N}_1)N_2 + N_1 \\ N_1^{(2)} = \text{hom}(\hat{N}_2, \hat{N}_1)N_2^{(1)} + N_1^{(1)}$$

as exhibited in the towers:

$$N_1^{(2)} = \text{hom}(\hat{N}_2, \hat{N}_1)N_2^{(1)} + N_1^{(1)} \quad N_2^{(2)} = \text{hom}(\hat{N}_1, \hat{N}_2)N_1^{(1)} + N_2^{(1)} \\ N_1^{(1)} = \text{hom}(\hat{N}_2, \hat{N}_1)N_2^{(0)} + N_1^{(0)} \quad N_2^{(1)} = \text{hom}(\hat{N}_1, \hat{N}_2)N_1^{(0)} + N_2^{(0)} \\ N_1^{(0)} = N_1 \quad N_2^{(0)} = N_2$$

Thus

$$\begin{aligned} N_1^{(2)} &= \text{hom}(\hat{N}_2, \hat{N}_1)[\text{hom}(\hat{N}_1, \hat{N}_2)N_1^{(0)} + N_2^{(0)}] + \text{hom}(\hat{N}_2, \hat{N}_1)N_2^{(0)} + N_1^{(0)} \\ &= \text{hom}(\hat{N}_2, \hat{N}_1) \text{hom}(\hat{N}_1, \hat{N}_2)N_1 + \text{hom}(\hat{N}_2, \hat{N}_1)N_2 + N_1. \end{aligned}$$

Next,

$$N_1^{(3)} = \text{hom}(\hat{N}_2, \hat{N}_1)N_2^{(2)} + N_1^{(2)}$$

where

$$N_2^{(2)} = \text{hom}(\hat{N}_1, \hat{N}_2)N_1^{(1)} + N_2^{(1)},$$

hence

$$N_2^{(2)} = \text{hom}(\hat{N}_1, \hat{N}_2) \text{hom}(\hat{N}_2, \hat{N}_1)N_2 + \text{hom}(\hat{N}_1, \hat{N}_2)N_1 + N_2.$$

Now

$$\text{hom}(\hat{N}_2, \hat{N}_1)N_2^{(2)} = \text{hom}(\hat{N}_2, \hat{N}_1)N_2 + \text{hom}(\hat{N}_2, \hat{N}_1) \text{hom}(\hat{N}_1, \hat{N}_2)N_1.$$

Adding  $N_1^{(2)}$  to both sides, we obtain

$$\begin{aligned} N_1^{(3)} &= N_1^{(2)} + \text{hom}(\hat{N}_2, \hat{N}_1)N_2 + \dots, \\ &= N_1 + \text{hom}(\hat{N}_2, \hat{N}_1)N_2 + \text{hom}(\hat{N}_2, \hat{N}_1) \text{hom}(\hat{N}_1, \hat{N}_2)N_1 = N_1^{(2)}. \end{aligned}$$

This yields by induction that  $N_i^{(k)} = N_i^{(2)} \forall k \geq 2$ , and hence  $\tilde{N}_i = N_i^{(2)}$ .

**EXAMPLE 3.3.** *It is possible to have a family  $\{N_j \mid j \in I\}$  where  $\tilde{N}_j \neq N_j^{(1)}$ .*

**PROOF.** Let  $V$  be an infinite dimensional left vector space over a field  $F$ . Let  $R = \text{End}_F(V)$ , and  $S = \text{Soc}(R_R) = \text{Soc}({}_R R)$ . It is well known that  ${}_R R$  is injective but  $R_R$  is not injective, and  $S_R$  is essential in  $R_R$ . Set  $M_1 = R$ ,  $M_2 = S$ . Then  $M_1^{(1)} = R$ ,  $M_2^{(1)} = E(R)$ , and  $\tilde{M}_1 = E(R) = \tilde{M}_2$ .

**4. Applications to quasi-injective and  $\pi$ -injective hulls.** Next we return to our consideration of a family of right  $R$ -modules  $\{M_i \mid i \in I\}$  where either  $I$  is finite or  $R$  is right noetherian and their direct sum  $M = \bigoplus_{i \in I} M_i$ . The mutual injective hulls discussed above in Theorems 3.1 and 3.2 allow us to develop formulae for  $q(M)$  and  $\pi(M)$ . But first,

**LEMMA 4.1.** *Let  $\{N_i \mid i \in I\}$  be a family of  $R$ -modules and  $f$  an endomorphism of  $E(N_i)$ . Then*

$$f(\tilde{N}_i) \subseteq \tilde{N}_i + f(N_i).$$

**PROOF.** Notice that  $N_i^{(k)} = \text{hom}(\sum_{j \neq i} N_j^{(k-1)}, \hat{N}_i) \sum_{j \neq i} N_j^{(k-1)} + N_i$ . Let  $g \in \text{hom}(\sum_{j \neq i} N_j^{(k-1)}, \hat{N}_i)$ . Then  $fg \in \text{hom}(\sum_{j \neq i} N_j^{(k-1)}, \hat{N}_i)$ . Hence  $f(N_i^{(k)}) \subseteq N_i^{(k)} + f(N_i)$ . Now the proof follows by the definition of  $\tilde{N}_i$ .

COROLLARY 4.2. *If  $N_i$  is quasi or  $\pi$ -injective, then so is  $\tilde{N}_i$ .*

THEOREM 4.3. *Let  $\{M_i \mid i \in I\}$  be a family of  $R$ -modules where either  $I$  is finite or  $R$  is right noetherian. Then*

$$\pi\left(\bigoplus_{i \in I} M_i\right) = \bigoplus_{i \in I} \pi(\tilde{M}_i) = \bigoplus_{i \in I} \pi(\widetilde{M}_i).$$

PROOF. Set  $\pi = \pi(\bigoplus_{i \in I} M_i)$ . Then  $E(\pi) = \bigoplus_{i \in I} E(M_i)$ , and so

$$\pi = \bigoplus_{i \in I} E(M_i) \cap \pi,$$

which yields  $\{E(M_i) \cap \pi \mid i \in I\}$  is mutually injective. Let  $\{\tilde{M}_i \mid i \in I\}$  be the mutual injective hull of  $\{M_i\}$  in  $\{E(M_i)\}$ . Then

$$\tilde{M}_i \subset \pi \cap E(M_i) \text{ for all } i.$$

Hence,  $\pi(\tilde{M}_i) \subset \pi$ ,  $\forall i$ . Thus  $\bigoplus \pi(\tilde{M}_i) \subset \pi$ . But  $\{\pi(\tilde{M}_i) \mid i \in I\}$  is mutually injective (Lemma 2.3). Hence,  $\sum \pi(\tilde{M}_i)$  is  $\pi$ -injective and equals  $\pi$ .

We next show the equality  $\pi = \bigoplus_{i \in I} \pi(\widetilde{M}_i)$ . Since  $\{\pi(\tilde{M}_i) \mid i \in I\}$  is a mutually injective family and for all  $i \in I$ ,  $\pi M_i \subseteq \pi \tilde{M}_i$ , we have by definition of mutual injective hull of this family  $\pi(\widetilde{M}_i) \subseteq \pi(\tilde{M}_i)$ , and so

$$(1) \quad \bigoplus_{i \in I} \pi(\widetilde{M}_i) \subseteq \bigoplus_{i \in I} \pi(\tilde{M}_i).$$

Next,

$$(2) \quad \begin{aligned} \bigoplus_{i \in I} \pi \tilde{M}_i &= \pi\left(\bigoplus_{i \in I} M_i\right) \subseteq \pi\left(\bigoplus_{i \in I} \pi M_i\right) \\ &= \bigoplus_{i \in I} \pi(\pi \tilde{M}_i) = \bigoplus_{i \in I} \pi(\widetilde{M}_i), \end{aligned}$$

by Corollary 4.2. Then by (1) and (2),  $\pi = \bigoplus_{i \in I} \pi(\widetilde{M}_i)$ , as claimed.

A similar result for quasi-injective hulls is contained in

THEOREM 4.4. *If  $I$  is finite or  $R$  is right noetherian, then*

$$q\left(\bigoplus_{i \in I} M_i\right) = \bigoplus_{i \in I} q(\tilde{M}_i) = \bigoplus_{i \in I} q(\widetilde{M}_i).$$

The first equality can actually be strengthened to obtain

$$q\left(\bigoplus_{i \in I} M_i\right) = \bigoplus_{i \in I} q(M_i^{(1)}).$$

PROOF. The proof of the first two equalities is identical to the proof of Theorem 4.3. For the latter equality, since

$$M_i^{(1)} = M_i + \sum_{j \neq i} \text{hom}(\hat{M}_j, \hat{M}_i) \text{hom}(\hat{M}_i, \hat{M}_j) M_i + \sum_{j \neq i} \text{hom}(\hat{M}_j, \hat{M}_i) M_j,$$

we get

$$qM_i^{(1)} = qM_i + \sum_{j \neq i} \text{hom}(\hat{M}_j, \hat{M}_i) \text{hom}(\hat{M}_i, \hat{M}_j) M_i + \sum_{j \neq i} \text{hom}(\hat{M}_j, \hat{M}_i) M_j,$$

which is clearly  $M_j$ -injective for all  $j \neq i$ , proving our claim.

EXAMPLE 4.5. *There exist modules  $M_1, M_2$  such that  $\pi(M_1 \oplus M_2) \neq \pi(M_1^{(1)}) \oplus \pi(M_2^{(1)})$ .*

PROOF. Let  $M_1 = R$  be a non-left artinian ring with right composition length 2 and essential simple right socle  $S = M_2$  ([2], p. 337). Then  $M_1^{(1)} = R, M_2^{(1)} = E(R)$ . So,  $\pi(M_1^{(1)}) \oplus \pi(M_2^{(1)}) = R \oplus E(R) \neq \pi(M_1 \oplus M_2) = E(R) \oplus E(R)$ .

ACKNOWLEDGEMENT. The authors are extremely grateful for the referee's valuable comments.

#### REFERENCES

1. F. W. Anderson and K. Fuller, *Rings and Categories of Modules*, Springer-Verlag, NY-Heidelberg, Berlin, (1993).
2. C. Faith, *Algebra I: Rings, Modules and Categories*, Springer-Verlag, New York, 1974.
3. V. K. Goel and S. K. Jain,  $\pi$ -injective modules and rings whose cyclic modules are  $\pi$ -injective, *Comm. Algebra* **6**(1978), 59–73.
4. S. H. Mohamed and B. J. Mueller, *Continuous and Discrete Modules*, London Math. Soc., LN 147, Cambridge University Press, (1990).
5. R. Wisbauer, *Foundations of Module and Ring Theory*, Gordon and Breach Sc. Pub., Philadelphia-Reading-Paris, (1991).

*Department of Mathematics*  
*Ohio University*  
*Athens, OH 45701*  
*U.S.A.*